

**POSTER PRESENTATION****Open Access**

Impact of gender on the prevalence and extent of microvascular obstruction after st-elevation myocardial infarction

Suzanne de Waha*, Ingo Eitel, Steffen Desch, Georg Fuernau, Philipp Lurz, Matthias Grothoff, Matthias Gutberlet, Gerhard Schuler, Holger Thiele

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Background

Baseline characteristics of women suffering from ST-elevation myocardial infarction (STEMI) diverge compared to men. Therefore the presence and extent of microvascular obstruction (MO), which has been shown to be a prognostic marker for adverse clinical outcome after STEMI, might differ between genders. Up to date gender differences of MO after STEMI have not been evaluated.

Methods

STEMI patients reperfused by primary angioplasty (n=423) within 12 hours after symptom onset underwent contrast-enhanced-MRI at a median of 3 (interquartile range [IQR] 2;4) days after the index event. MO was measured 15 minutes after gadolinium injection with late enhancement sequences and evaluated qualitatively and quantitatively (as percentage of left ventricular mass [%LV]).

Results

A total of 105 women and 318 men were analysed (24.6 vs. 76.4%, $p < 0.001$). In comparison to men, women were significantly older (71.5 [IQR 63.4;77.0] vs. 63.7 [IQR 54.0;71.0] years, $p < 0.001$), displayed longer symptom-onset-to-reperfusion times (241 [IQR 156;389] vs. 196 [IQR 132;337] minutes, $p = 0.04$), a higher prevalence of diabetes mellitus (35.9 vs. 23.3%, $p = 0.02$) and arterial hypertension (76.7 vs. 65.6%, $p = 0.04$). Complete epicardial reperfusion defined as post-PCI TIMI-flow III (95.2 vs. 95.2%, $p = 1.0$) did not differ significantly between genders.

The prevalence and extent of MO, infarct size and left ventricular ejection fraction were similar in women and

men (MO presence: 65.4 vs. 71.9%, $p = 0.22$ / MO extent: 0.59 [IQR 0;1.24] vs. 0.71 [IQR 0;2.0] %LV, $p = 0.47$; infarct size: 13.9 [IQR 5.8;25.6] vs. 18.4 [IQR 8.7;29.0] % LV, $p = 0.14$; ejection fraction: 52.2 [IQR 42.6;60.0] vs. 49.4 [40.7;57.9] %, $p = 0.16$).

Conclusion

Despite longer ischemic time and a more disadvantageous cardiovascular risk profile in women the prevalence and extent of MO do not differ between genders.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P103

Cite this article as: de Waha et al.: Impact of gender on the prevalence and extent of microvascular obstruction after st-elevation myocardial infarction. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1): P103.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



University of Leipzig – Heart Center, Leipzig, Germany
Full list of author information is available at the end of the article



© 2011 de Waha et al; licensee BioMed Central Ltd. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.